

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HER-2 / CD340(HRB2/451), CF405S conjugate, 0.1mg/mL, Clone: [HRB2/451], Mouse, Monoclonal BOT-BNC040451-100

| | |
|--------------------------|---|
| Artikelname | HER-2 / CD340(HRB2/451), CF405S conjugate, 0.1mg/mL, Clone: [HRB2/451], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040451-100 |
| Hersteller Artikelnummer | BNC040451-100 |
| Alternativnummer | BOT-BNC040451-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human HER-2 protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb is specific to c-erbB-2/HER-2 and shows minimal cross-reaction with other members of the family. C-erbB-2/HER-2 is a member of the EGFR family. Receptors of this family are located on the plasma membrane and consist of an extracellular ligan... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [HRB2/451] |

| | |
|------------------------|--|
| Molekulargewicht | 185 kDa |
| UniProt | P04626 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |