

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin 1 / EMA / Episialin / CD227(VU-2G7), CF405S conjugate, 0.1mg/mL, Clone: [VU-2G7], Mouse, Monoclonal BOT-BNC040630-500

| | |
|--------------------------|---|
| Artikelname | Mucin 1 / EMA / Episialin / CD227(VU-2G7), CF405S conjugate, 0.1mg/mL, Clone: [VU-2G7], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040630-500 |
| Hersteller Artikelnummer | BNC040630-500 |
| Alternativnummer | BOT-BNC040630-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Synthetic glycosylated MUC1 60mer tandem repeat NH2-(HGVTSAPDT(GalNAc)RPAPGSTAPPAHG)3- COOH, conjugated to bovine serum albumin |
| Konjugation | CF405S |
| Produktbeschreibung | MAb VU-2G7 reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of VU-2G7 has not been established with epitope fingerprinting. The extracellular domain of MUC1 large... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [VU-2G7] |
| Molekulargewicht | 265-400 kDa |
| UniProt | P15941 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-paraffin 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |