

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Myogenin(F5D), CF405S conjugate, 0.1mg/mL, Clone: [F5D], Mouse, Monoclonal BOT-BNC040632-500**

Artikelname	Myogenin(F5D), CF405S conjugate, 0.1mg/mL, Clone: [F5D], Mouse, Monoclonal
Artikelnummer	BOT-BNC040632-500
Hersteller Artikelnummer	BNC040632-500
Alternativnummer	BOT-BNC040632-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Canine, Feline, Human, Mouse, Porcine, Rat
Immunogen	Rat myogenin peptide (aa 73-94) followed by rat myogenin recombinant fragment (aa30-224) (Epitope aa138-158)
Konjugation	CF405S
Produktbeschreibung	Myogenin is a member of the MyoD family of myogenic basic helix-loop-helix (bHLH) transcription factors that also includes MyoD, Myf-5, and MRF4 (also known as herculinor Myf-6). MyoD family members are expressed exclusively in skeletal muscle and pl...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[F5D]
Molekulargewicht	34 kDa
UniProt	<a href="#">P15173</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user