

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **von Willebrand Factor(3E2D10), CF405S conjugate, 0.1mg/mL, Clone: [3E2D10], Mouse, Monoclonal BOT-BNC040633-100**

Artikelname	von Willebrand Factor(3E2D10), CF405S conjugate, 0.1mg/mL, Clone: [3E2D10], Mouse, Monoclonal
Artikelnummer	BOT-BNC040633-100
Hersteller Artikelnummer	BNC040633-100
Alternativnummer	BOT-BNC040633-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human vWF fragment spanning aa 845-949
Konjugation	CF405S
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[3E2D10]

Molekulargewicht	250 kDa
UniProt	<a href="#">P04275</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user