

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD31 / PECAM-1(JC70/A + C31.3), CF405S conjugate, 0.1mg/mL, Clone: [JC70/A C31.3], Mouse, Monoclonal BOT-BNC040647-500

| | |
|--------------------------|---|
| Artikelname | CD31 / PECAM-1(JC70/A + C31.3), CF405S conjugate, 0.1mg/mL, Clone: [JC70/A C31.3], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040647-500 |
| Hersteller Artikelnummer | BNC040647-500 |
| Alternativnummer | BOT-BNC040647-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human, Primate, Rabbit |
| Immunogen | Human recombinant CD31 protein (C31.3) & Membrane preparation of a spleen from a patient with hairy cell leukemia (JC/70A) |
| Konjugation | CF405S |
| Produktbeschreibung | CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [JC70/A C31.3] |
| Molekulargewicht | ~100 kDa (endothelium) and ~130 kDa (platelets) |
| UniProt | P16284 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |