

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TNF alpha(TNF655), CF405S conjugate, 0.1mg/mL, Clone: [TNF655], Mouse, Monoclonal BOT-BNC040655-500**

|                          |   |
|--------------------------|---|
| Artikelname              | TNF alpha(TNF655), CF405S conjugate, 0.1mg/mL, Clone: [TNF655],<br>Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC040655-500   |
| Hersteller Artikelnummer | BNC040655-500   |
| Alternativnummer         | BOT-BNC040655-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human TNF-alpha   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | This MAb recognizes human 17-26 kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17 kDa free molecule, or as a 26 kDa membrane protein. TNF-alpha is a protein secreted by lipopolysa... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [TNF655]  |
| Molekulargewicht         | 17 kDa  |

|                        |  |
|------------------------|--|
| UniProt                | <a href="#">P01375</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |