

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27 / KIP1(SX53G8), CF405S conjugate, 0.1mg/mL, Clone: [SX53G8], Mouse, Monoclonal BOT-BNC040669-100

| | |
|--------------------------|---|
| Artikelname | p27 / KIP1(SX53G8), CF405S conjugate, 0.1mg/mL, Clone: [SX53G8], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040669-100 |
| Hersteller Artikelnummer | BNC040669-100 |
| Alternativnummer | BOT-BNC040669-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, IP, WB |
| Spezies Reaktivität | Human, Monkey, Mouse, Rat |
| Immunogen | Purified GST-p27 fusion protein of human origin |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb recognizes a 27 kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human brea... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [SX53G8] |

| | |
|------------------------|--|
| Molekulargewicht | 25-26 kDa |
| UniProt | P46527 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |