

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Hepatocyte Specific(OCH1E5), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [OCH1E5], Mouse, Monoclonal BOT-BNC040671-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Hepatocyte Specific(OCH1E5), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [OCH1E5], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC040671-500   |
| Hersteller Artikelnummer | BNC040671-500   |
| Alternativnummer         | BOT-BNC040671-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Canine, Human   |
| Immunogen                | Extract of a formalin-fixed, rejected-allograft of a human liver  |
| Konjugation              | CF405S  |
| Produktbeschreibung      | Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or HepPar1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [OCH1E5]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | Not Known   |
| Isotyp                 | IgG1  |
| UniProt                | Not Known   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |