

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

pS2(GE2), CF405S conjugate, 0.1mg/mL, Clone: [GE2], Mouse, Monoclonal BOT-BNC040677-500

| | |
|--------------------------|---|
| Artikelname | pS2(GE2), CF405S conjugate, 0.1mg/mL, Clone: [GE2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040677-500 |
| Hersteller Artikelnummer | BNC040677-500 |
| Alternativnummer | BOT-BNC040677-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human pS2. |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes a polypeptide of 6.5 kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [GE2] |
| Molekulargewicht | 6.5 kDa |
| UniProt | P04155 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |