

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD34(HPCA1/763), CF405S conjugate, 0.1mg/mL, Clone: [HPCA1/763], Mouse, Monoclonal BOT-BNC040763-100

| | |
|--------------------------|---|
| Artikelname | CD34(HPCA1/763), CF405S conjugate, 0.1mg/mL, Clone: [HPCA1/763], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040763-100 |
| Hersteller Artikelnummer | BNC040763-100 |
| Alternativnummer | BOT-BNC040763-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human HPCA1 protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120 kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually l... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [HPCA1/763] |

| | |
|------------------------|---|
| Molekulargewicht | 90-110 kDa |
| UniProt | P28906 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |