

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **IgA Secretory Component(ECM1/792), CF405S conjugate, 0.1mg/mL, Clone: [ECM1/792], Mouse, Monoclonal BOT-BNC040792-500**

Artikelname	IgA Secretory Component(ECM1/792), CF405S conjugate, 0.1mg/mL, Clone: [ECM1/792], Mouse, Monoclonal
Artikelnummer	BOT-BNC040792-500
Hersteller Artikelnummer	BNC040792-500
Alternativnummer	BOT-BNC040792-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human ECM1 protein
Konjugation	CF405S
Produktbeschreibung	This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[ECM1/792]

Molekulargewicht	~80 kDa
UniProt	<a href="#">Q16610</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>