

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Retinol Binding Protein-1(G4E4), CF405S conjugate, 0.1mg/mL, Clone: [G4E4], Mouse, Monoclonal BOT-BNC040855-100

| | |
|--------------------------|---|
| Artikelname | Retinol Binding Protein-1(G4E4), CF405S conjugate, 0.1mg/mL, Clone: [G4E4], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040855-100 |
| Hersteller Artikelnummer | BNC040855-100 |
| Alternativnummer | BOT-BNC040855-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Goat, Human, Monkey, Mouse, Primate, Rabbit, Rat |
| Immunogen | Retinol binding protein-1 purified from human plasma |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb recognizes an epitope within the 74-182 C-terminal sequence (11kD peptide fragment) of human serum Cellular Retinol Binding Protein 1 (CRBP 1), a single-chain glycoprotein belonging to the superfamily of hydrophobic molecule transporter prot... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [G4E4] |

| | |
|------------------------|---|
| Molekulargewicht | 21-25 kDa |
| UniProt | P02753 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |