

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glypican-3(GPC3/863), CF405S conjugate, 0.1mg/mL, Clone: [GPC3/863], Mouse, Monoclonal BOT-BNC040863-100

| | |
|--------------------------|--|
| Artikelname | Glypican-3(GPC3/863), CF405S conjugate, 0.1mg/mL, Clone: [GPC3/863], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040863-100 |
| Hersteller Artikelnummer | BNC040863-100 |
| Alternativnummer | BOT-BNC040863-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human GPC3 protein |
| Konjugation | CF405S |
| Produktbeschreibung | Glypican-3 (GPC3) is a glycosylphosphatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, mel... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [GPC3/863] |

| | |
|------------------------|---|
| Molekulargewicht | 67 kDa |
| UniProt | P51654 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |