

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD99 / MIC2 (Ewings Sarcoma Marker)(12E7 + MIC2/877), CF405S conjugate, 0.1mg/mL, Clone: [12E7 MIC2/877], Mouse, Monoclonal BOT-BNC040878-100

Artikelname	CD99 / MIC2 (Ewings Sarcoma Marker)(12E7 + MIC2/877), CF405S conjugate, 0.1mg/mL, Clone: [12E7 MIC2/877], Mouse, Monoclonal
Artikelnummer	BOT-BNC040878-100
Hersteller Artikelnummer	BNC040878-100
Alternativnummer	BOT-BNC040878-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human acute lymphocytic leukemia T-cells (12E7), Recombinant human MIC2 protein (MIC2/877)
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 27-32 kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which ar...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[12E7 MIC2/877]
Molekulargewicht	27-32 kDa
UniProt	P14209
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Flow cytometry: 5-10 uL/million cells Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1:100-1:200 for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Immunofluorescence 1:100-1:200 Optimal dilution for a specific application should be determined by user