

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

GFAP(GA-5 + ASTRO/789), CF405S conjugate, 0.1mg/mL, Clone: [GA-5 ASTRO/789], Mouse, Monoclonal BOT-BNC040898-500

| | |
|--------------------------|---|
| Artikelname | GFAP(GA-5 + ASTRO/789), CF405S conjugate, 0.1mg/mL, Clone: [GA-5 ASTRO/789], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040898-500 |
| Hersteller Artikelnummer | BNC040898-500 |
| Alternativnummer | BOT-BNC040898-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Bovine, Gallus, Human, Mouse, Porcine, Rabbit, Rat |
| Immunogen | GFAP isolated from pig spinal cord (GA-5), Recombinant GFAP protein (ASTRO/789) |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb recognizes a protein of ~50 kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for I... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [GA-5 ASTRO/789] |
| Molekulargewicht | ~50 kDa |
| UniProt | P14136 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |