

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Androgen Receptor(AR441 + DHTR/882), CF405S conjugate, 0.1mg/mL, Clone: [AR441 DHTR/882], Mouse, Monoclonal BOT-BNC040913-500

| | |
|--------------------------|---|
| Artikelname | Androgen Receptor(AR441 + DHTR/882), CF405S conjugate, 0.1mg/mL, Clone: [AR441 DHTR/882], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040913-500 |
| Hersteller Artikelnummer | BNC040913-500 |
| Alternativnummer | BOT-BNC040913-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | A synthetic peptide, aa 299-315, (STEDTAEYSPFKGGYTK) of human AR (AR441), Recombinant human DHTR protein (DHTR/882) |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes a protein of 110 kDa, which is identified as androgen receptor (AR). It reacts with full length, and the newly described A form of the receptor. It does not cross react with estrogen, progesterone, or glucocorticoid receptors... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [AR441 DHTR/882] |
| Molekulargewicht | ~110 kDa |
| UniProt | P10275 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Does not react with mouse, others not known Optimal dilution for a specific application should be determined by user |