

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin 6 (MUC6)(MUC6/916), CF405S conjugate, 0.1mg/mL, Clone: [MUC6/916], Mouse, Monoclonal BOT-BNC040916-500

| | |
|--------------------------|---|
| Artikelname | Mucin 6 (MUC6)(MUC6/916), CF405S conjugate, 0.1mg/mL, Clone: [MUC6/916], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040916-500 |
| Hersteller Artikelnummer | BNC040916-500 |
| Alternativnummer | BOT-BNC040916-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human MUC6 protein |
| Konjugation | CF405S |
| Produktbeschreibung | The MUC6 gastric mucin is a secreted glycoprotein that plays an essential role in epithelial cyto-protection from acid, proteases, pathogenic microorganisms, and mechanical trauma in the gastrointestinal tract. Mucin 6 expression is highest in the st... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [MUC6/916] |

| | |
|------------------------|---|
| Molekulargewicht | 252 kDa |
| UniProt | Q6W4X9 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |