

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myosin, Smooth Muscle Heavy Chain(MYH11/923), CF405S conjugate, 0.1mg/mL, Clone: [MYH11/923], Mouse, Monoclonal BOT-BNC040923-500

| | |
|--------------------------|---|
| Artikelname | Myosin, Smooth Muscle Heavy Chain(MYH11/923), CF405S conjugate, 0.1mg/mL, Clone: [MYH11/923], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040923-500 |
| Hersteller Artikelnummer | BNC040923-500 |
| Alternativnummer | BOT-BNC040923-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human MYH11 protein |
| Konjugation | CF405S |
| Produktbeschreibung | Smooth muscle myosin heavy chain (SM-MHC) is a cytoplasmic structural protein, which is a major component of the contractile apparatus in smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early in smooth ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [MYH11/923] |

| | |
|------------------------|---|
| Molekulargewicht | 205 kDa (MHC-1) and 200 kDa (MHC-2) |
| UniProt | P35749 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Predicted to show broad species reactivity Optimal dilution for a specific application should be determined by user |