

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

bcl-2(BCL2/782 + BCL2/796), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal BOT-BNC041023-500

Artikelname	bcl-2(BCL2/782 + BCL2/796), CF405S conjugate, 0.1mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal
Artikelnummer	BOT-BNC041023-500
Hersteller Artikelnummer	BNC041023-500
Alternativnummer	BOT-BNC041023-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human bcl-2 protein
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[BCL2/782 BCL2/796]

Molekulargewicht	25-26 kDa
UniProt	P10415
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user