

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SOX10(SOX10/991 + SOX10/1074), CF405S conjugate, 0.1mg/mL, Clone: [SOX10/991 SOX10/1074], Mouse, Monoclonal BOT-BNC041074-100

| | |
|--------------------------|---|
| Artikelname | SOX10(SOX10/991 + SOX10/1074), CF405S conjugate, 0.1mg/mL, Clone: [SOX10/991 SOX10/1074], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041074-100 |
| Hersteller Artikelnummer | BNC041074-100 |
| Alternativnummer | BOT-BNC041074-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (aa 115-269) of human SOX10 protein with hexa-histidine tag |
| Konjugation | CF405S |
| Produktbeschreibung | Recognizes a protein of ~55 kDa, identified as SOX10. This MAb is highly specific and does not cross-react with other members of the SOX-family. SOX genes comprise a family of genes that are related to the mammalian sex-determining gene SRY. These ge... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [SOX10/991 SOX10/1074] |
| Molekulargewicht | 49-58 kDa |
| UniProt | P56693 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user</p> |