

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD63 (Late Endosomes Marker)(NK1/C3 + LAMP3/968), CF405S conjugate, 0.1mg/mL, Clone: [NK1/C3 LAMP3/968], Mouse, Monoclonal BOT-BNC041160-100**

Artikelname	CD63 (Late Endosomes Marker)(NK1/C3 + LAMP3/968), CF405S conjugate, 0.1mg/mL, Clone: [NK1/C3 LAMP3/968], Mouse, Monoclonal
Artikelnummer	BOT-BNC041160-100
Hersteller Artikelnummer	BNC041160-100
Alternativnummer	BOT-BNC041160-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Smooth plasma membrane fraction of MeWo cells (NK1/C3), Recombinant human full-length CD63 protein (LAMP3/968)
Konjugation	CF405S
Produktbeschreibung	This MAb recognizes protein of 26 kDa-60 kDa, which is identified as CD63. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with sp...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NKI/C3 LAMP3/968]
Molekulargewicht	26 kDa (core protein), 30-60 kDa (glycosylated)
UniProt	<a href="#">P08962</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user