

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PTH / Parathyroid Hormone (N-Terminal)(PTH/1173), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1173], Mouse, Monoclonal BOT-BNC041173-500

| | |
|--------------------------|---|
| Artikelname | PTH / Parathyroid Hormone (N-Terminal)(PTH/1173), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1173], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041173-500 |
| Hersteller Artikelnummer | BNC041173-500 |
| Alternativnummer | BOT-BNC041173-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (84 amino acid residues from C-terminus) of human PTH protein |
| Konjugation | CF405S |
| Produktbeschreibung | The epitope of this MAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood calcium levels by dissolving the salts i... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [PTH/1173] |
| Molekulargewicht | 9 kDa |
| UniProt | P01270 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |