

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **EMI1(EMI1/1176), CF405S conjugate, 0.1mg/mL, Clone: [EMI1/1176], Mouse, Monoclonal BOT-BNC041176-100**

Artikelname	EMI1(EMI1/1176), CF405S conjugate, 0.1mg/mL, Clone: [EMI1/1176], Mouse, Monoclonal
Artikelnummer	BOT-BNC041176-100
Hersteller Artikelnummer	BNC041176-100
Alternativnummer	BOT-BNC041176-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (203 amino acid residues between aa 1-250) of human EMI1 protein
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a 56 kDa protein, which is identified as Early Mitotic Inhibitor-1 (EMI1). It regulates mitosis by inhibiting the anaphase promoting complex/cyclosome (APC). Emi1 is a conserved F box protein containing a zinc-binding region ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[EMI1/1176]
Molekulargewicht	56 kDa
UniProt	<a href="#">Q9UKT4</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user