

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Topoisomerase II alpha (Proliferation & Drug-Resistance Marker) (TOP2A/1361), CF405S conjugate, 0.1mg/mL, Clone: [TOP2A/1361], Mouse, Monoclonal BOT-BNC041361-500**

Artikelname	Topoisomerase II alpha (Proliferation & Drug-Resistance Marker) (TOP2A/1361), CF405S conjugate, 0.1mg/mL, Clone: [TOP2A/1361], Mouse, Monoclonal
Artikelnummer	BOT-BNC041361-500
Hersteller Artikelnummer	BNC041361-500
Alternativnummer	BOT-BNC041361-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant human Topoisomerase II alpha fragment (aa1352-1493) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a 170 kDa protein, which is identified as topoisomerase IIalpha. It shows no cross-reaction with topoisomerase IIbeta or topoisomerase I. Topo IIalpha plays important roles in synthesis and transcription of DNA as well as chr...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[TOP2A/1361]
Molekulargewicht	170 kDa
UniProt	<a href="#">P11388</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user