

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Lewis Y (Tumor Marker) (LWY/1463), CF405S conjugate, 0.1mg/mL, Clone: [LWY/1463], Mouse, Monoclonal BOT-BNC041463-100

| | |
|--------------------------|---|
| Artikelname | Lewis Y (Tumor Marker) (LWY/1463), CF405S conjugate, 0.1mg/mL, Clone: [LWY/1463], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041463-100 |
| Hersteller Artikelnummer | BNC041463-100 |
| Alternativnummer | BOT-BNC041463-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human colon carcinoma cells |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucalpha1-2Galbeta1-4/3GlcNAcbeta). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classification of human ren... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [LWY/1463] |

| | |
|------------------------|---|
| Molekulargewicht | Multiple |
| UniProt | Not Applicable |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |