

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), CF405S conjugate, 0.1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal BOT-BNC041496-100

| | |
|--------------------------|---|
| Artikelname | CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), CF405S conjugate, 0.1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041496-100 |
| Hersteller Artikelnummer | BNC041496-100 |
| Alternativnummer | BOT-BNC041496-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human NCAM1 protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [NCAM1/1496] |

| | |
|------------------------|--|
| Molekulargewicht | 125 kDa 145 180 |
| UniProt | P13591 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |