

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TNF-alpha (Tumor Necrosis Factor alpha) (TNF/1500R), CF405S conjugate, 0.1mg/mL, Clone: [TNF/1500R], Rabbit, Monoclonal BOT-BNC041500-500

| | |
|--------------------------|---|
| Artikelname | TNF-alpha (Tumor Necrosis Factor alpha) (TNF/1500R), CF405S conjugate, 0.1mg/mL, Clone: [TNF/1500R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC041500-500 |
| Hersteller Artikelnummer | BNC041500-500 |
| Alternativnummer | BOT-BNC041500-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human TNF-alpha protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb recognizes human 17-26 kDa protein, which is identified as cytokine TNF-alpha (Tumor Necrosis Factor-alpha). TNF-alpha can be expressed as a 17 kDa free molecule, or as a 26 kDa membrane protein. TNF-alpha is a protein secreted by lipopolysa... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TNF/1500R] |

| | |
|------------------------|--|
| Molekulargewicht | 17 kDa |
| UniProt | P01375 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 4-8 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |