

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Ep-CAM / CD326 (Epithelial Marker), CF405S conjugate, 0.1mg/mL, Rabbit BOT-BNC041647-500

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Ep-CAM / CD326 (Epithelial Marker), CF405S conjugate, 0.1mg/mL, Rabbit                                                                                                                                                                                        |
| Artikelnummer            | BOT-BNC041647-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC041647-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC041647-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human TACSTD1 protein                                                                                                                                                                                                                             |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo... |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Molekulargewicht         | 40-43 kDa                                                                                                                                                                                                                                                     |
| UniProt                  | <a href="#">P16422</a>                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |