

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100A (Melanoma Marker) (S1/61), CF405S conjugate, 0.1mg/mL, Clone: [S1/61], Mouse, Monoclonal BOT-BNC041695-100

| | |
|--------------------------|---|
| Artikelname | S100A (Melanoma Marker) (S1/61), CF405S conjugate, 0.1mg/mL, Clone: [S1/61], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041695-100 |
| Hersteller Artikelnummer | BNC041695-100 |
| Alternativnummer | BOT-BNC041695-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified human S-100 protein conjugated to methylated BSA |
| Konjugation | CF405S |
| Produktbeschreibung | S100A and S100B proteins are two members of the S100 family of calcium binding proteins. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the alpha-ch... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [S1/61] |

| | |
|------------------------|---|
| Molekulargewicht | 10-12 kDa |
| UniProt | P23297 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |