

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CTLA4 / CD152 (Negative Regulator of T-Cells)(L4P2F5.F10), CF405S conjugate, 0.1mg/mL, Clone: [L4P2F5.F10], Mouse, Monoclonal BOT-BNC041696-100**

Artikelname	CTLA4 / CD152 (Negative Regulator of T-Cells)(L4P2F5.F10), CF405S conjugate, 0.1mg/mL, Clone: [L4P2F5.F10], Mouse, Monoclonal
Artikelnummer	BOT-BNC041696-100
Hersteller Artikelnummer	BNC041696-100
Alternativnummer	BOT-BNC041696-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	A synthetic peptide from CTLA4 (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	CTLA4 (CD152) is a cell surface receptor that behaves as a negative regulator of the proliferation and the effector function of T cells. It contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[L4P2F5.F10]
Molekulargewicht	30-43 kDa

UniProt	<a href="#">P16410</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry