

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TCF4 (Transcription Factor 4) (TCF4/1705), CF405S conjugate, 0.1mg/mL, Clone: [TCF4/1705], Mouse, Monoclonal BOT-BNC041705-500**

|                          |   |
|--------------------------|---|
| Artikelname              | TCF4 (Transcription Factor 4) (TCF4/1705), CF405S conjugate, 0.1mg/mL, Clone: [TCF4/1705], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC041705-500   |
| Hersteller Artikelnummer | BNC041705-500   |
| Alternativnummer         | BOT-BNC041705-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | WB  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human TCF4 protein fragment (around aa 365-671) (exact sequence is proprietary)   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | This antibody recognizes a protein of 71 kDa, identified as Transcription Factor 4 (TCF4). It is a basic helix-loop-helix transcription factor. The protein recognizes an Ephrussi-box (E-box) binding site (CANNTG) - a motif first identified in immunog... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [TCF4/1705]   |
| Molekulargewicht       | 71 kDa  |
| UniProt                | <a href="#">P15884</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | ELISA: Use Ab at 2-4 ug/mL for coating, Order Ab without BSA, Western blot 1-2 ug/mL, Optimal dilution for a specific application should be determined.   Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody |