

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Parathyroid Hormone (PTH) (N-Terminal) (PTH/1717R), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1717R], Rabbit, Monoclonal BOT-BNC041717-100

| | |
|--------------------------|---|
| Artikelname | Parathyroid Hormone (PTH) (N-Terminal) (PTH/1717R), CF405S conjugate, 0.1mg/mL, Clone: [PTH/1717R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC041717-100 |
| Hersteller Artikelnummer | BNC041717-100 |
| Alternativnummer | BOT-BNC041717-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human PTH protein N-terminal fragment (exact sequence is proprietary) |
| Konjugation | CF405S |
| Produktbeschreibung | Epitope of this MAb maps in the N-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood calcium levels by dissolving the salts in bo... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [PTH/1717R] |
| Molekulargewicht | 9 kDa |
| UniProt | P01270 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |