

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase (Melanoma Marker) (TYR/2024R), CF405S conjugate, 0.1mg/mL, Clone: [TYR/2024R], Rabbit, Monoclonal BOT-BNC042024-100

| | |
|--------------------------|---|
| Artikelname | Tyrosinase (Melanoma Marker) (TYR/2024R), CF405S conjugate, 0.1mg/mL, Clone: [TYR/2024R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC042024-100 |
| Hersteller Artikelnummer | BNC042024-100 |
| Alternativnummer | BOT-BNC042024-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human tyrosinase (TYR) protein |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes a cluster of proteins between 70-80 kDa, identified as tyrosinase. Occasionally a minor band at 55 kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TYR/2024R] |

| | |
|------------------------|--|
| Molekulargewicht | 70-80 kDa |
| UniProt | P14679 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 or 10 mM citrate buffer pH6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |