

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), CF405S conjugate, 0.1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal BOT-BNC042034-100

Artikelname	CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/2034R), CF405S conjugate, 0.1mg/mL, Clone: [LPFS2/2034R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042034-100
Hersteller Artikelnummer	BNC042034-100
Alternativnummer	BOT-BNC042034-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD22 protein (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[LPFS2/2034R]
Molekulargewicht	120 kDa
UniProt	P26842
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user