

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EpCAM / CD326 (Epithelial Marker) (EGP40/2041R), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/2041R], Rabbit, Monoclonal BOT-BNC042041-500

Artikelname	EpCAM / CD326 (Epithelial Marker) (EGP40/2041R), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/2041R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042041-500
Hersteller Artikelnummer	BNC042041-500
Alternativnummer	BOT-BNC042041-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human EpCAM fragment from extracellular domain (around aa100-224) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	Epitope of this MAb is mapped between aa 202-212 of EGP40, which is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-la...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[EGP40/2041R]
Molekulargewicht	40-43 kDa
UniProt	P16422
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user