

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD43 (T-Cell Marker) (SPN/2049R), CF405S conjugate, 0.1mg/mL, Clone: [SPN/2049R], Rabbit, Monoclonal BOT-BNC042049-500**

Artikelname	CD43 (T-Cell Marker) (SPN/2049R), CF405S conjugate, 0.1mg/mL, Clone: [SPN/2049R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042049-500
Hersteller Artikelnummer	BNC042049-500
Alternativnummer	BOT-BNC042049-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD43 protein
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43. From 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[SPN/2049R]

Molekulargewicht	95, 115, or 135 kDa
UniProt	<a href="#">P16150</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user