

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nucleophosmin (Acute Myeloid Leukemia Marker) (NA24), CF405S conjugate, 0.1mg/mL, Clone: [NA24], Mouse, Monoclonal BOT-BNC042053-100

Artikelname	Nucleophosmin (Acute Myeloid Leukemia Marker) (NA24), CF405S conjugate, 0.1mg/mL, Clone: [NA24], Mouse, Monoclonal
Artikelnummer	BOT-BNC042053-100
Hersteller Artikelnummer	BNC042053-100
Alternativnummer	BOT-BNC042053-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	GST fusion protein containing the N-terminal part of nucleophosmin fused to 14 amino acid of ALK protein
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a 33 kDa glycoprotein, identified as Nucleophosmin (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is high...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NA24]
Molekulargewicht	33 kDa
UniProt	P06748
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user