

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD7 (T-Cell Leukemia Marker)(CD7/3737), CF405S conjugate, 0.1mg/mL, Clone: [CD7/3737], Mouse, Monoclonal BOT-BNC042056-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD7 (T-Cell Leukemia Marker)(CD7/3737), CF405S conjugate, 0.1mg/mL, Clone: [CD7/3737], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC042056-100   |
| Hersteller Artikelnummer | BNC042056-100   |
| Alternativnummer         | BOT-BNC042056-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human full-length CD7 protein   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | Recognizes a protein of 40 kDa, identified as CD7 (also known as gp40, Leu9). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [CD7/3737]  |
| Molekulargewicht         | 40 kDa  |

|                        |  |
|------------------------|--|
| UniProt                | <a href="#">P09564</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |