

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TCL1 (T-Cell Marker) (TCL1/2078), CF405S conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal BOT-BNC042078-100

Artikelname	TCL1 (T-Cell Marker) (TCL1/2078), CF405S conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal
Artikelnummer	BOT-BNC042078-100
Hersteller Artikelnummer	BNC042078-100
Alternativnummer	BOT-BNC042078-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[TCL1/2078]
Molekulargewicht	14 kDa
UniProt	P56279
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user