

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53 Tumor Suppressor Protein (rTP53/1739), CF405S conjugate, 0.1mg/mL, Clone: [rTP53/1739], Mouse, Monoclonal BOT-BNC042091-100

| | |
|--------------------------|---|
| Artikelname | p53 Tumor Suppressor Protein (rTP53/1739), CF405S conjugate, 0.1mg/mL, Clone: [rTP53/1739], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC042091-100 |
| Hersteller Artikelnummer | BNC042091-100 |
| Alternativnummer | BOT-BNC042091-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human TP53 protein |
| Konjugation | CF405S |
| Produktbeschreibung | The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM) Array, containing more than 19,000, full-length human proteins. Recognizes a 53 kDa protein, which is identified as p53 suppressor gene product. It reacts ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [rTP53/1739] |

| | |
|------------------------|---|
| Molekulargewicht | 53 kDa |
| UniProt | P04637 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |