

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53 Tumor Suppressor Protein (rBP53-12), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal BOT-BNC042094-100

| | |
|--------------------------|---|
| Artikelname | p53 Tumor Suppressor Protein (rBP53-12), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC042094-100 |
| Hersteller Artikelnummer | BNC042094-100 |
| Alternativnummer | BOT-BNC042094-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Canine, Gallus, Hamster, Human, Monkey |
| Immunogen | Recombinant human wild-type p53 protein |
| Konjugation | CF405S |
| Produktbeschreibung | This MAb reacts with an N-terminal epitope (aa 16-25) of both wild type and mutated p53. Mutation and/or allelic loss of p53 is one of the causes of a variety of mesenchymal and epithelial tumors. If it occurs in the germ line, such tumors run in fam... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [rBP53-12] |

| | |
|------------------------|--|
| Molekulargewicht | 53 kDa |
| Isotyp | IgG1 |
| UniProt | P04637 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |