

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p53 Tumor Suppressor Protein (rBP53-12), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal BOT-BNC042094-500**

Artikelname	p53 Tumor Suppressor Protein (rBP53-12), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rBP53-12], Mouse, Monoclonal
Artikelnummer	BOT-BNC042094-500
Hersteller Artikelnummer	BNC042094-500
Alternativnummer	BOT-BNC042094-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Canine, Gallus, Hamster, Human, Monkey
Immunogen	Recombinant human wild-type p53 protein
Konjugation	CF405S
Produktbeschreibung	This MAb reacts with an N-terminal epitope (aa 16-25) of both wild type and mutated p53. Mutation and/or allelic loss of p53 is one of the causes of a variety of mesenchymal and epithelial tumors. If it occurs in the germ line, such tumors run in fam...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rBP53-12]

Molekulargewicht	53 kDa
Isotyp	IgG1
UniProt	<a href="#">P04637</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user