

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), CF405S conjugate, 0.1mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal BOT-BNC042153-100

Artikelname	TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), CF405S conjugate, 0.1mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal
Artikelnummer	BOT-BNC042153-100
Hersteller Artikelnummer	BNC042153-100
Alternativnummer	BOT-BNC042153-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[TACSTD2/2153]
Molekulargewicht	40 kDa
UniProt	P09758
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user