

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8 (KRT8) (KRT8/2174R), CF405S conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal BOT-BNC042174-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 8 (KRT8) (KRT8/2174R), CF405S conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal                                                                                                                                                        |
| Artikelnummer            | BOT-BNC042174-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC042174-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC042174-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC, WB                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Bovine, Human, Porcine, Rabbit, Sheep                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human KRT8 protein                                                                                                                                                                                                                    |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [KRT8/2174R]                                                                                                                                                                                                                                                  |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 52.5 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| UniProt                | <a href="#">P05787</a>                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |