

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF405S conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal BOT-BNC042289-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Human Kappa Light Chain / IGKC (B-Cell Marker) (KLC2289R), CF405S conjugate, 0.1mg/mL, Clone: [KLC2289R], Rabbit, Monoclonal  |
| Artikelnummer            | BOT-BNC042289-500   |
| Hersteller Artikelnummer | BNC042289-500   |
| Alternativnummer         | BOT-BNC042289-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human Ig kappa chain (IGKC) protein   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. It recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects al... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [KLC2289R]  |
| Molekulargewicht       | ~22.5 kDa   |
| UniProt                | <a href="#">P01601</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |