

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **STAT3 / Signal Transducer and Activator of Transcription 3 (STAT3/2409), CF405S conjugate, 0.1mg/mL, IgG2b, Clone: [STAT3/2409], Mouse, Monoclonal BOT-BNC042409-100**

Artikelname	STAT3 / Signal Transducer and Activator of Transcription 3 (STAT3/2409), CF405S conjugate, 0.1mg/mL, IgG2b, Clone: [STAT3/2409], Mouse, Monoclonal
Artikelnummer	BOT-BNC042409-100
Hersteller Artikelnummer	BNC042409-100
Alternativnummer	BOT-BNC042409-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human STAT3 protein
Konjugation	CF405S
Produktbeschreibung	The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM) Array, containing more than 19,000, full-length human proteins. STAT3 is a member of the STAT protein family. In response to cytokines and growth factors, ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[STAT3/2409]
Molekulargewicht	88 kDa
Isotyp	IgG2b
UniProt	<a href="#">P40763</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user