

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), CF405S conjugate, 0.1mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal BOT-BNC042498-100

| | |
|--------------------------|---|
| Artikelname | Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), CF405S conjugate, 0.1mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC042498-100 |
| Hersteller Artikelnummer | BNC042498-100 |
| Alternativnummer | BOT-BNC042498-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary) |
| Konjugation | CF405S |
| Produktbeschreibung | Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different core... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [rBCL6/1718] |
| Molekulargewicht | 95 kDa |
| UniProt | P41182 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |