

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD44 / HCAM Std. (HCAM/2875R), CF405S conjugate, 0.1mg/mL, Clone: [HCAM/2875R], Rabbit, Monoclonal BOT-BNC042875-100**

Artikelname	CD44 / HCAM Std. (HCAM/2875R), CF405S conjugate, 0.1mg/mL, Clone: [HCAM/2875R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042875-100
Hersteller Artikelnummer	BNC042875-100
Alternativnummer	BOT-BNC042875-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes
Konjugation	CF405S
Produktbeschreibung	This antibody recognizes a cell surface glycoprotein of 80-95 kDa (CD44) on lymphocytes, monocytes, and granulocytes (Leucocyte Typing Workshop V). Its epitope is resistant to digestion by trypsin and chymotrypsin. This MAb selectively interferes wit...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[HCAM/2875R]

Molekulargewicht	80-95 kDa
UniProt	<a href="#">P16070</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user