

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD6 (Negative Marker of T-regulatory Cells) (C6/2884R), CF405S conjugate, 0.1mg/mL, Clone: [C6/2884R], Rabbit, Monoclonal BOT-BNC042884-500

Artikelname	CD6 (Negative Marker of T-regulatory Cells) (C6/2884R), CF405S conjugate, 0.1mg/mL, Clone: [C6/2884R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042884-500
Hersteller Artikelnummer	BNC042884-500
Alternativnummer	BOT-BNC042884-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Human recombinant full-length CD6 protein
Konjugation	CF405S
Produktbeschreibung	CD6 is a type I transmembrane glycoprotein that contains a 24-amino acid signal sequence, three extracellular scavenger receptor cysteine-rich (SRCR) domains, a membrane-spanning domain and a 44-amino acid cytoplasmic domain. The CD6 glycoprotein is ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[C6/2884R]

Molekulargewicht	90-130 kDa
UniProt	P30203
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user