

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Secretory Component / ECM1 (ECM1/2889R), CF405S conjugate, 0.1mg/mL, Clone: [ECM1/2889R], Rabbit, Monoclonal BOT-BNC042889-100**

Artikelname	Secretory Component / ECM1 (ECM1/2889R), CF405S conjugate, 0.1mg/mL, Clone: [ECM1/2889R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042889-100
Hersteller Artikelnummer	BNC042889-100
Alternativnummer	BOT-BNC042889-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human ECM1 protein
Konjugation	CF405S
Produktbeschreibung	This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[ECM1/2889R]

Molekulargewicht	~80 kDa
UniProt	<a href="#">Q16610</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user